

WHAT IS CLAIMED IS:

1. A method of identifying a candidate beta catenin pathway modulating agent,
5 said method comprising the steps of:
 - (a) providing an assay system comprising a UP polypeptide or nucleic acid;
 - (b) contacting the assay system with a test agent under conditions whereby,
but for the presence of the test agent, the system provides a reference activity; and
 - (c) detecting a test agent-biased activity of the assay system, wherein a
10 difference between the test agent-biased activity and the reference activity identifies the
test agent as a candidate beta catenin pathway modulating agent.
2. The method of Claim 1 wherein the assay system comprises cultured cells that
express the UP polypeptide.
15
3. The method of Claim 2 wherein the cultured cells additionally have defective
beta catenin function.
4. The method of Claim 1 wherein the assay system includes a screening assay
20 comprising a UP polypeptide, and the candidate test agent is a small molecule modulator.
5. The method of Claim 4 wherein the assay is a phosphorylase assay.
6. The method of Claim 1 wherein the assay system is selected from the group
25 consisting of an apoptosis assay system, a cell proliferation assay system, an angiogenesis
assay system, and a hypoxic induction assay system.
7. The method of Claim 1 wherein the assay system includes a binding assay
comprising a UP polypeptide and the candidate test agent is an antibody.
30
8. The method of Claim 1 wherein the assay system includes an expression assay
comprising a UP nucleic acid and the candidate test agent is a nucleic acid modulator.

9. The method of claim 8 wherein the nucleic acid modulator is an antisense oligomer.

10. The method of Claim 8 wherein the nucleic acid modulator is a PMO.

5

11. The method of Claim 1 additionally comprising:

(d) administering the candidate beta catenin pathway modulating agent identified in (c) to a model system comprising cells defective in beta catenin function and, detecting a phenotypic change in the model system that indicates that the beta catenin function is restored.

10

12. The method of Claim 11 wherein the model system is a mouse model with defective beta catenin function.

15

13. A method for modulating a beta catenin pathway of a cell comprising contacting a cell defective in beta catenin function with a candidate modulator that specifically binds to a UP polypeptide, whereby beta catenin function is restored.

20

14. The method of claim 13 wherein the candidate modulator is administered to a vertebrate animal predetermined to have a disease or disorder resulting from a defect in beta catenin function.

25

15. The method of Claim 13 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.

16. The method of Claim 1, comprising the additional steps of:

(d) providing a secondary assay system comprising cultured cells or a non-human animal expressing UP ,

(e) contacting the secondary assay system with the test agent of (b) or an agent derived therefrom under conditions whereby, but for the presence of the test agent or agent derived therefrom, the system provides a reference activity; and

30

(f) detecting an agent-biased activity of the second assay system,

wherein a difference between the agent-biased activity and the reference activity of the second assay system confirms the test agent or agent derived therefrom as a candidate beta catenin pathway modulating agent,

5 and wherein the second assay detects an agent-biased change in the beta catenin pathway.

17. The method of Claim 16 wherein the secondary assay system comprises cultured cells.

10 18. The method of Claim 16 wherein the secondary assay system comprises a non-human animal.

19. The method of Claim 18 wherein the non-human animal mis-expresses a beta catenin pathway gene.

15 20. A method of modulating beta catenin pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a UP polypeptide or nucleic acid.

20 21. The method of Claim 20 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the beta catenin pathway.

22. The method of Claim 20 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.

25 23. A method for diagnosing a disease in a patient comprising:
obtaining a biological sample from the patient;
contacting the sample with a probe for UP expression;
comparing results from step (b) with a control;
30 determining whether step (c) indicates a likelihood of disease.

24. The method of claim 23 wherein said disease is cancer.

25. The method according to claim 24, wherein said cancer is a cancer as shown in
35 Table 1 as having >25% expression level.